

SJSU/TP-99-20
June 1999

Quantum Probability from Decision Theory?

J. Finkelstein¹
Department of Physics
San José State University
San José, CA 95192, U.S.A

Abstract

Deutsch has recently (in quant-ph/9906015) offered a justification, based only on the non-probabilistic axioms of quantum theory and of classical decision theory, for the use of the standard quantum probability rules. In this note, this justification is examined.

¹ Participating Guest, Lawrence Berkeley National Laboratory
e-mail: JLFINKELSTEIN@lbl.gov

The origin, and indeed the meaning, of the probability rules of quantum theory are often considered mysterious. In a recent article (ref. [1], hereinafter denote by [D]), David Deutsch has offered a proof, based only on the non-probabilistic axioms of quantum theory and of classical decision theory, that a “rational decision maker” would act as if the standard probability rules of quantum theory were true. (See also ref. [2].) This result could then be taken to justify these standard rules, without the need to introduce any probabilistic axiom, or even the need to have any definition of probability. It is the purpose of this note to examine the proof given in [D].

[D] considers a decision maker who is offered the following game: an observable $\hat{X}$ is to be measured on a quantum system which is in the state $|\Psi\rangle$; the utility of the payoff offered is numerically equal to the result of the measurement. The value of this game is denoted by $\mathcal{V}[|\Psi\rangle]$, and the proof given in [D] is that, even without any probabilistic axioms, it follows that

$$\mathcal{V}[|\Psi\rangle] = \langle\Psi|\hat{X}|\Psi\rangle. \quad (1)$$

The RHS of Eq. 1 is of course the value of the game that would follow from the standard quantum probability rules; the derivation of Eq. 1 without invoking any probability axioms can thus be considered a justification of those rules.

I wish to examine a central part of the proof offered in [D], in which the state $|\Psi\rangle$ is an equal-amplitude superposition of two eigenstates of $\hat{X}$. I will use the notation employed in [D], that x is an eigenvalue, and $|x\rangle$ the corresponding eigenstate, of the operator $\hat{X}$. Then the states we will be concerned with can be written

$$|\Psi\rangle = \frac{1}{\sqrt{2}}(|x_1\rangle + |x_2\rangle); \quad (2)$$

this is also Eq. 7 of [D]. The non-probabilistic part of quantum theory implies that the outcome of the measurement of $\hat{X}$ on the state given in Eq. 2 is surely either x_1 or x_2 (but of course does not specify which one); from this fact, together with axioms of classical decision theory, [D] deduces that (Eq. 11 in [D])

$$\mathcal{V}[\frac{1}{\sqrt{2}}(|x_1\rangle + |x_2\rangle)] = \frac{1}{2}(x_1 + x_2). \quad (3)$$

This result (our Eq. 3) is called the “pivotal result” of [D].

But how could this result follow from the argument made in [D]? Let r denote the outcome of the measurement of $\hat{X}$ on the state given in Eq. 2; then the only requirement of quantum theory used in the proof of Eq. 3 is that

either $r = x_1$ or $r = x_2$. Now suppose (consistent with that requirement) that the value of r were in fact determinate, and were given (in the case $|x_1| \neq |x_2|$) by

$$\begin{aligned} r &= x_1 & \text{if } |x_1| > |x_2| \\ r &= x_2 & \text{if } |x_2| > |x_1|. \end{aligned} \quad (4)$$

When the outcome of the measurement is determinate, the value of the game is simply equal to the outcome, so if, for example, we set $x_1 = 1$ and $x_2 = 0$, this supposition gives $r = 1$, and so the LHS of Eq. 3 would be 1 while the RHS of that equation would be 0.5.

The proof of Eq. 3 (Eq. 11 of [D]) begins with Eq. 8 of [D]. That equation, specialized to the case of an equal-amplitude superposition of two eigenstates, reads

$$\mathcal{V}[\frac{1}{\sqrt{2}}(|x_1 + k\rangle + |x_2 + k\rangle)] = k + \mathcal{V}[\frac{1}{\sqrt{2}}(|x_1\rangle + |x_2\rangle)], \quad (5)$$

and is supposed to be true for any k . However, if the notation in Eq. 5 is taken to mean the same as that in Eqs. 1 and 2 (e.g., that $|x_1 + k\rangle$ is the eigenstate corresponding to eigenvalue $x_1 + k$), the game on the LHS of Eq. 5 may not even be possible. In writing Eq. 2, we have assumed that the operator $\hat{X}$ has eigenvalues x_1 and x_2 , but since nothing else was assumed about the spectrum of this operator, it might not have eigenvalues $x_1 + k$ and $x_2 + k$, and so the state required by the game on the LHS of Eq. 5 might not exist. Furthermore, in the case in which this state *does* happen to exist, the argument given in [D] would allow this equation to be incorrect. Consider the example given in Eq. 4, and take $x_1 = 1$, $x_2 = 0$, and $k = -1$. Then since, by Eq. 4, $\mathcal{V}[\frac{1}{\sqrt{2}}(|0\rangle + |-1\rangle)] = -1$ and $\mathcal{V}[\frac{1}{\sqrt{2}}(|1\rangle + |0\rangle)] = +1$, Eq. 5 reads $-1 = 0$.

On the other hand, it is possible to re-interpret the notation used in Eq. 5 in a way that makes it correct. Let me set $|x_1, x_2\rangle =: \frac{1}{\sqrt{2}}(|x_1\rangle + |x_2\rangle)$, and define

$$\mathcal{V}[|x_1, x_2\rangle; x_1 \rightarrow u_1, x_2 \rightarrow u_2]$$

to be the value of the following game: $\hat{X}$ is measured on the state $|x_1, x_2\rangle$; if the outcome is x_1 , the utility is u_1 , while if the outcome is x_2 , the utility is u_2 . The relation between this notation and the original notation is

$$\mathcal{V}[\frac{1}{\sqrt{2}}(|x_1\rangle + |x_2\rangle)] = \mathcal{V}[|x_1, x_2\rangle; x_1 \rightarrow x_1, x_2 \rightarrow x_2]. \quad (6)$$

Now if we interpret the LHS of Eq. 5 not as $\mathcal{V}[|x_1 + k, x_2 + k\rangle; x_1 + k \rightarrow x_1 + k, x_2 + k \rightarrow x_2 + k]$, but rather as $\mathcal{V}[|x_1, x_2\rangle; x_1 \rightarrow x_1 + k, x_2 \rightarrow x_2 + k]$,

we can re-write Eq. 5 as

$$\mathcal{V}[|x_1, x_2\rangle; x_1 \rightarrow x_1 + k, x_2 \rightarrow x_2 + k] = k + \mathcal{V}[|x_1, x_2\rangle; x_1 \rightarrow x_1, x_2 \rightarrow x_2]. \quad (7)$$

This equation is meaningful (because the same state $|x_1, x_2\rangle$ is required in the two games), and moreover it is correct. So perhaps this is the way we were intended to understand Eq. 8 of [D].

We can now attempt to complete the derivation of Eq. 3, along the line followed in [D]. Setting $k = -x_1 - x_2$ in Eq. 7, we get

$$\mathcal{V}[|x_1, x_2\rangle; x_1 \rightarrow -x_2, x_2 \rightarrow -x_1] = -x_1 - x_2 + \mathcal{V}[|x_1, x_2\rangle; x_1 \rightarrow x_1, x_2 \rightarrow x_2]. \quad (8)$$

In this notation, the “zero-sum rule” (Eq. 9 of [D]) reads

$$\mathcal{V}[|x_1, x_2\rangle; x_1 \rightarrow u_1, x_2 \rightarrow u_2] + \mathcal{V}[|x_1, x_2\rangle; x_1 \rightarrow -u_1, x_2 \rightarrow -u_2] = 0, \quad (9)$$

and if we use this (with $u_1 = -x_2$ and $u_2 = -x_1$) in Eq. 8, we get

$$-\mathcal{V}[|x_1, x_2\rangle; x_1 \rightarrow x_2, x_2 \rightarrow x_1] = -x_1 - x_2 + \mathcal{V}[|x_1, x_2\rangle; x_1 \rightarrow x_1, x_2 \rightarrow x_2]. \quad (10)$$

This is our version of Eq. 10 of [D]. However, we cannot now go from this equation to Eq. 3 (i.e., to Eq. 11 of [D]), since the values of the two games which appear in Eq. 10 have not been shown to be equal (Of course, they *are* equal under the condition that the two outcomes x_1 and x_2 are equally probable, but that is precisely the condition that [D] does *not* want to assume). In fact, in the example given by Eq. 4 with $x_1 = 1$ and $x_2 = 0$, the outcome is surely x_1 , and so

$$\begin{aligned} \mathcal{V}[|x_1, x_2\rangle; x_1 \rightarrow x_1, x_2 \rightarrow x_2] &= x_1 = 1 \\ \mathcal{V}[|x_1, x_2\rangle; x_1 \rightarrow x_2, x_2 \rightarrow x_1] &= x_2 = 0; \end{aligned} \quad (11)$$

Eq. 10 is satisfied (both sides equal 0), but Eq. 3 is not.

We are certainly not suggesting that Eq. 3 and the more general Eq. 1 (i.e., Eqs. 11 and 4 of [D]) are not correct; they do follow, after all, from the usual quantum probability rules. Furthermore, we are not claiming to have shown that these equations are not consequences of just the assumptions (the non-probabilistic parts of quantum theory and of decision theory) that were made in [D]. We do see, however, that the arguments given in [D] are not sufficient to establish the validity of these equations.

Acknowledgement: I would like to acknowledge the hospitality of the Lawrence Berkeley National Laboratory, where this work was done.

References

- [1] D. Deutsch, “Quantum Theory of Probability and Decisions,” preprint quant-ph/9906015, to appear in Proc. R. Soc. Lond. **A**
- [2] L. Polley, “Reducing the Statistical Axiom,” preprint quant-ph/9906124.